Primes In Arithmetic Progressions And Primitive Roots

N. A. Carella

***Abstract*:** Let $x \geq 1$ be a large number, and let $1 \leq a < q$ be integers such that $\gcd(a,q) = 1$ and $q = O(\log^c)$ with $c > 0$ constant. This note proves that the counting function for the number of primes $p \in \{p = qn + a : n \geq 1\}$ with a fixed primitive root $u \neq \pm 1, v^2$ has the asymptotic formula $\pi_u(x,q,a) = \delta(u,q,a)x/\log x + O(x/\log^b x)$, where $\delta(u,q,a) > 0$ is the density, and $b > c+1$ is a constant.

Mathematics Subjects Classification: 11A07, 11N37.
Keywords: Prime Number; Primitive Root; Arithmetic Progression.

arXiv:1701.03188v1 [math.GM] 11 Jan 2017

1 Introduction

Let u, q and a be integers with $\gcd(a,q) = 1$. The density of the subset of primes in the arithmetic progression $\{p = qn + a : \gcd(a,q) = 1 \text{ and } \operatorname{ord}_p(u) = p-1\}$ is defined by $\delta(u,q,a)$, and the corresponding counting function

$$\pi_u(x,q,a) = \#\{p = qn + a \leq x : \gcd(a,q) = 1 \text{ and } \operatorname{ord}_p(u) = p-1\} \tag{1}$$

for any large number $x \geq 1$ are subject of this note.

Theorem 1.1 *Let $x \geq 1$ be a large number. Let $1 \leq a < q$ be integers such that $\gcd(a,q) = 1$ and $q = O(\log^c x)$, with $c \geq 0$ constant. Then, the arithmetic progression $\{p = qn + a : n \geq 1\}$ has infinitely many primes $p \geq 2$ with a fixed primitive root $u \neq \pm 1, v^2$. In addition, the counting function for these primes has the asymptotic formula*

$$\pi_u(x,q,a) = \delta(u,q,a)\frac{x}{\log x} + O\left(\frac{x}{\log^b x}\right), \tag{2}$$

where $\delta(u,q,a) \geq 0$ is the density constant depending on the integers u, q, a, and $b > c+1$ is a constant.

The preliminary background results and notation are discussed in Sections two to five. Section six presents a proof of Theorem 1.1. Lastly, Corollary 7.1 provides the density for the Gauss problem about the primitive root $u = 10$.

2 Representation of the Characteristic Function

The *order* of an element in the cyclic group $G = \mathbb{F}_p^\times$ is defined by

$$\operatorname{ord}_p(v) = \min\{k : v^k \equiv 1 \bmod p\}. \tag{3}$$

Primitive elements in this cyclic group have order $p - 1 = \#G$. The characteristic function $\Psi : G \longrightarrow \{0,1\}$ of primitive elements is one of the standard analytic tools employed to investigate the various properties of primitive roots in cyclic groups G. Many equivalent representations of the characteristic function Ψ of primitive elements are possible. The standard characteristic function is discussed in [11, p. 258]. It detects a primitive element by means of the divisors of $p - 1$.

A new representation of the characteristic function for primitive elements is developed here. It detects the order $\mathrm{ord}_p(u) \geq 1$ of the element $u \in \mathbb{F}_p$ by means of the solutions of the equation $\tau^n - u = 0$ in $\mathbb{F}_p$, where u, τ are constants, and n is a variable such that $1 \leq n < p-1, \gcd(n, p-1) = 1$.

Lemma 2.1 *Let $p \geq 2$ be a prime, and let τ be a primitive root mod p. Let $u \in \mathbb{F}_p$ be a nonzero element, and let, $\psi \neq 1$ be a nonprincipal additive character of order* $\mathrm{ord}\,\psi = p$. *Then*

$$\Psi(u) = \sum_{\gcd(n,p-1)=1} \frac{1}{p} \sum_{0 \leq k \leq p-1} \psi\left((\tau^n - u)k\right) = \begin{cases} 1 & \text{if } \mathrm{ord}_p(u) = p-1, \\ 0 & \text{if } \mathrm{ord}_p(u) \neq p-1. \end{cases} \tag{4}$$

Proof: As the index $n \geq 1$ ranges over the integers relatively prime to $p-1$, the element $\tau^n \in \mathbb{F}_p$ ranges over the primitive roots $\mathrm{mod}\, p$. Ergo, the equation $\tau^n - u = 0$ has a solution if and only if the fixed element $u \in \mathbb{F}_p$ is a primitive root. Next, replace $\psi(z) = e^{i2\pi kz/p}$ to obtain

$$\sum_{\gcd(n,p-1)=1} \frac{1}{p} \sum_{0 \leq k \leq p-1} e^{i2\pi(\tau^n - u)k/p} = \begin{cases} 1 & \text{if } \mathrm{ord}_p(u) = p-1, \\ 0 & \text{if } \mathrm{ord}_p(u) \neq p-1. \end{cases} \tag{5}$$

This follows from the geometric series identity $\sum_{0 \leq n \leq N-1} w^n = (w^N - 1)/(w-1), w \neq 1$ applied to the inner sum. ■.

3 Estimates Of Exponential Sums

This Section provides a simple estimate for the exponential sum of interest in this analysis. The proof of this estimate is entirely based on next theorem and elementary techniques. Some versions of the next result for double exponential sums are also proved in [2], and [5].

Theorem 3.1 *([6, Theorem 1]) Let $p \geq 2$ be a large prime, and let $\tau \in \mathbb{F}_p$ be an element of large multiplicative order $T = \mathrm{ord}_p(\tau)$. Let $X, Y \subset \mathbb{F}_p$ be large subsets of cardinality $\#X \geq \#Y \geq p^{7/8+\varepsilon}$ Then, for any given number $k > 0$, the following estimate holds.*

$$\sum_{y \in Y} \left| \sum_{x \in X} \gamma(x) e^{i2\pi a \tau^{xy}/p} \right| \ll \frac{\#Y^{1-\frac{1}{2k}} \cdot \#X^{1-\frac{1}{2k+2}} \cdot p^{\frac{1}{2k} + \frac{3}{4k+4} + o(1)}}{T^{\frac{1}{2k+2}}}, \tag{6}$$

where the coefficients $\gamma(x)$ are complex numbers, $|\gamma(x)| \leq 1$, and $a \geq 1, \gcd(a,p) = 1$ is an integer.

Lemma 3.1 *Let $p \geq 2$ be a large prime, and let τ be a primitive root modulo p. Then,*

$$\max_{\gcd(s,p-1)=1} \left| \sum_{\gcd(n,p-1)=1} e^{i2\pi s \tau^n/p} \right| \ll p^{1-\varepsilon} \tag{7}$$

for any small number $\varepsilon < 1/16$.

Proof: Let $X = Y = \{n : \gcd(n, p-1) = 1\} \subset \mathbb{F}_p$ be subsets of cardinalities $\#X = \#Y = \varphi(p-1)$. Since for each fixed $y \in Y$, the map $x \longrightarrow xy$ is a permutation of the subset $X \subset \mathbb{F}_p$, the double sum can be rewritten as a single exponential sum

$$\sum_{y \in Y} \left| \sum_{x \in X} \gamma(x) e^{i2\pi s \tau^{xy}/p} \right| = \#Y \left| \sum_{x \in X} e^{i2\pi s \tau^x/p} \right|, \tag{8}$$

where $\gamma(x) = 1$ if $x \in X$ otherwise it vanishes. As the element τ is a primitive root, it has order $T = \mathrm{ord}_p(\tau) = p-1$. Setting $k = 1$ in Theorem 3.1 yields

$$\begin{aligned}\left|\sum_{x\in X} e^{i2\pi s\tau^x/p}\right| &= \frac{1}{\#Y}\sum_{y\in Y}\left|\sum_{x\in X} e^{i2\pi s\tau^{xy}/p}\right| \\ &\ll \frac{1}{\#Y}\left(\#Y^{1/2}\cdot \#X^{3/4}\cdot \frac{p^{7/8+\varepsilon}}{T^{1/4}}\right) \\ &\ll \frac{1}{\varphi(p-1)}\left(\varphi(p-1)^{1/2}\cdot\varphi(p-1)^{3/4}\cdot\frac{p^{7/8+\varepsilon}}{(p-1)^{1/4}}\right) \\ &\ll p^{7/8+\varepsilon}.\end{aligned} \tag{9}$$

This is restated in the simpler notation $p^{7/8+\varepsilon} \le p^{1-\varepsilon}$ for any number $\varepsilon \in (0, 1/16)$. ■.

A similar result appears as Theorem 6 in [3].

4 Estimate For The Error Term

The upper bound for exponential sum over subsets of elements in finite fields $\mathbb{F}_p$ stated in the last Section will be used here to estimate the error term $E(x)$ arising in the proof of Theorem 1.1.

Lemma 4.1 *Let $p \ge 2$ be a large prime, let $\psi \ne 1$ be an additive character, and let τ be a primitive root mod p. If the element $u \ne 0$ is not a primitive root, then,*

$$\left|\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}}\frac{1}{p}\sum_{\gcd(n,p-1)=1,\ 0<k\le p-1}\psi\left((\tau^n-u)k\right)\right| \ll \frac{1}{\varphi(q)}\frac{x^{1-\varepsilon}}{\log x}, \tag{10}$$

where $1 \le a < q$, $\gcd(a,q) = 1$ and $O(\log^c x)$ with $c > 0$ constant, for all sufficiently large numbers $x \ge 1$ and an arbitrarily small number $\varepsilon > 0$.

Proof: By hypothesis $u \ne \tau^n$ for any $n \ge 1$ such that $\gcd(n, p-1) = 1$. Therefore, $\sum_{0<k\le p-1}\psi\left((\tau^n-u)k\right) = -1$, and

$$\begin{aligned}|E(x)| &= \left|\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}}\frac{1}{p}\sum_{\gcd(n,p-1)=1,\ 0<k\le p-1}\psi\left((\tau^n-u)k\right)\right| \\ &\le \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}}\frac{\varphi(p-1)}{p} \\ &\le \frac{1}{2}\frac{1}{\varphi(q)}\frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right),\end{aligned} \tag{11}$$

where $\varphi(n)/n \le 1/2$ for any integer $n \ge 1$. This implies that there is a nontrivial upper bound. To sharpen this upper bound, let $\psi(z) = e^{i2\pi kz/p}$ with $0 < k < p$, and rearrange the triple finite sum in the form

$$\begin{aligned}|E(x)| &= \left|\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}}\frac{1}{p}\sum_{0<k\le p-1,\ \gcd(n,p-1)=1}\psi((\tau^n-u)k)\right| \\ &\le \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}}\left|\frac{1}{p}\sum_{0<k\le p-1}e^{-i2\pi uk/p}\sum_{\gcd(n,p-1)=1}e^{i2\pi k\tau^n/p}\right|,\end{aligned} \tag{12}$$

and let

$$U_p = \frac{1}{p^{1/2}} \sum_{0<k\le p-1} e^{-i2\pi uk/p} \qquad \text{and} \qquad V_p = \frac{1}{p^{1/2}} \sum_{\gcd(n,p-1)=1} e^{i2\pi k\tau^n/p}. \tag{13}$$

Now consider the Holder inequality $||AB||_1 \le ||A||_r \cdot ||B||_s$ with $1/r + 1/s = 1$. In terms of the components in (13) this inequality has the explicit form

$$\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |U_pV_p| \le \left(\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |U_p|^r \right)^{1/r} \left(\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |V_p|^s \right)^{1/s}. \tag{14}$$

The absolute value of the first exponential sum U_p is given by

$$|U_p| = \left| \frac{1}{p^{1/2}} \sum_{0<k\le p-1} e^{-i2\pi uk/p} \right| = \frac{1}{p^{1/2}}. \tag{15}$$

This follows from $\sum_{0<k\le p-1} e^{i2\pi uk/p} = -1$ for $u \ne 0$ and summation of the geometric series. The corresponding r-norm has the upper bound

$$\begin{aligned} ||U_p||_r^r &= \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |U_p|^r \\ &= \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} \left| \frac{1}{p^{1/2}} \right|^r \\ &\le \frac{1}{x^{r/2}} \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} 1 \\ &\le \frac{3}{\varphi(q)} \frac{x^{1-r/2}}{\log x}. \end{aligned} \tag{16}$$

Here the finite sum over the primes is estimated using the Brun-Titchmarsh theorem; this result states that the number of primes $p = qn + 1$ in the interval $[x, 2x]$ satisfies the inequality

$$\pi(2x, q, a) - \pi(x, q, a) \le \frac{3}{\varphi(q)} \frac{x}{\log x}, \tag{17}$$

see [10, p. 167], [8, p. 157], [12], and [20, p. 83].

The absolute value of the second exponential sum $V_p = V_p(k)$ has the upper bound

$$|V_p| = \left| \frac{1}{p^{1/2}} \sum_{\gcd(n,p-1)=1} e^{i2\pi k\tau^n} \right| \ll p^{1/2-\varepsilon}. \tag{18}$$

This exponential sum dependents on k; but it has a uniform, and independent of k upper bound

$$\max_{1\le k\le p-1} \left| \sum_{\gcd(n,p-1)=1,} e^{i2\pi k\tau^n/p} \right| \ll p^{1-\varepsilon}, \tag{19}$$

where $\varepsilon > 0$ is an arbitrarily small number, see Lemma **??**. A similar application appears in [15, p. 1286].

In light of this estimate, the corresponding s-norm has the upper bound

$$\begin{aligned} ||V_p||_s^s &= \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |V_p|^s \\ &\ll \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} \left|p^{1/2-\varepsilon}\right|^s \\ &\ll \left((2x)^{1/2-\varepsilon}\right)^s \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} 1 \\ &\ll \frac{1}{\varphi(q)}\frac{(2x)^{1+s/2-\varepsilon s}}{\log x} \\ &\ll \frac{1}{\varphi(q)}\frac{x^{1+s/2-\varepsilon s}}{\log x}. \end{aligned} \tag{20}$$

The finite sum over the primes in the third line is estimated using the Brun-Titchmarsh theorem; and the last line uses $2^{1/s+1/2-\varepsilon} \le 2$ for $s \ge 1$.

Now, replace the estimates (16) and (20) into (14), the Holder inequality, to reach

$$\begin{aligned} \sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |U_pV_p| &\le \left(\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |U_p|^r\right)^{1/r} \left(\sum_{\substack{x\le p\le 2x\\ p\equiv a \bmod q}} |V_p|^s\right)^{1/s} \\ &\ll \left(\frac{1}{\varphi(q)}\frac{x^{1-r/2}}{\log x}\right)^{1/r} \left(\frac{1}{\varphi(q)}\frac{x^{1+s/2-\varepsilon s}}{\log x}\right)^{1/s} \\ &\ll \frac{1}{\varphi(q)}\frac{x^{1-\varepsilon}}{\log x} \\ &\ll \frac{1}{\varphi(q)}\frac{x^{1-\varepsilon}}{\log x}. \end{aligned} \tag{21}$$

Note that this result is independent of the parameter $1/r + 1/s = 1$. ■.

5 Evaluation Of The Main Term

Finite sums and products over the primes numbers occur on various problems concerned with primitive roots. These sums and products often involve the normalized totient function $\varphi(n)/n = \prod_{p|n}(1-1/p)$ and the corresponding estimates, and the asymptotic formulas.

Lemma 5.1 *([14, Lemma 5]) Let $x \ge 1$ be a large number, and let $\varphi(n)$ be the Euler totient function. If $q \le \log^c x$, with $c \ge 0$ constant, an integer $1 \le a < q$ such that $\gcd(a,q) = 1$, then*

$$\sum_{\substack{p\le x\\ p\equiv a \bmod q}} \frac{\varphi(p-1)}{p-1} = A_q\frac{\mathrm{li}(x)}{\varphi(q)} + O\left(\frac{x}{\log^B x}\right), \tag{22}$$

where $\mathrm{li}(x)$ is the logarithm integral, and $b > c+1 \ge 1$ is an arbitrary constant, as $x \to \infty$, and

$$A_q = \prod_{p|\gcd(a-1,q)} \left(1-\frac{1}{p}\right) \prod_{p\nmid q} \left(1-\frac{1}{p(p-1)}\right). \tag{23}$$

Related discussions for $q = 2$ are given in [19, Lemma 1], [13, p. 16], and, [21]. The case $q = 2$ is ubiquitous in various results in Number Theory.

Lemma 5.2 *Let $x \geq 1$ be a large number, and let $\varphi(n)$ be the Euler totient function. If $q \leq \log^c x$, with $c \geq 0$ constant, an integer $1 \leq a < q$ such that $\gcd(a, q) = 1$, then*

$$\sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{1}{p} \sum_{\gcd(n, p-1)=1} 1 = A_q \frac{\mathrm{li}(x)}{\varphi(q)} + O\left(\frac{x}{\log^b x}\right), \tag{24}$$

where $\mathrm{li}(x)$ is the logarithm integral, and $b > c + 1 \geq 1$ is an arbitrary constant, as $x \to \infty$, and A_q is defined in (23).

Proof: A routine rearrangement gives

$$\begin{aligned} \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{1}{p} \sum_{\gcd(n, p-1)=1} 1 &= \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p} \\ &= \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p-1} - \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p(p-1)}. \end{aligned} \tag{25}$$

To proceed, apply Lemma 5.1 to reach

$$\begin{aligned} \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p-1} - \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p(p-1)} &= A_q \frac{\mathrm{li}(x)}{\varphi(q)} + O\left(\frac{x}{\log^b x}\right) \\ &\quad - \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p(p-1)} \\ &= A_q \frac{\mathrm{li}(x)}{\varphi(q)} + O\left(\frac{x}{\log^b x}\right), \end{aligned} \tag{26}$$

where the second finite sum

$$\sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \frac{\varphi(p-1)}{p(p-1)} \ll \log\log x \tag{27}$$

is absorbed into the error term, $b > c + 1 \geq 1$ is an arbitrary constant, and A_q is defined in (23). ■.
The logarithm integral has the asymptotic formula

$$\mathrm{li}(x) = \int_2^x \frac{1}{\log z} dz = \frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right). \tag{28}$$

6 The Main Result

Given a fixed integer $u \neq \pm 1, v^2$, the precise primes counting function is defined by

$$\pi_u(x, q, a) = \#\{p \leq x : p \equiv a \bmod q \text{ and } \mathrm{ord}_p(u) = p - 1\} \tag{29}$$

for $1 \leq a < q$ and $\gcd(a, q) = 1$. The limit

$$\delta(u, q, a) = \lim_{x \to \infty} \frac{\pi_\ell(x, q, a)}{\pi(x, q, a)} = a_u \frac{A_q}{\varphi(q)} \tag{30}$$

is the density of the subset of primes with a fixed primitive root $u \neq \pm 1, v^2$. The proof of Theorem 1.1 is completed here.

Proof: Suppose that $u \neq \pm 1, v^2$ is not a primitive root for all primes $p \geq x_0$, with $x_0 \geq 1$ constant. Let $x > x_0$ be a large number, and $q = O(\log^c x)$. Consider the sum of the characteristic function over the short interval $[x, 2x]$, that is,

$$0 = \sum_{\substack{x \leq p \leq 2x \\ q \equiv a \bmod q}} \Psi(u). \tag{31}$$

Replacing the characteristic function, Lemma **??**, and expanding the nonexistence equation (31) yield

$$\begin{aligned} 0 &= \sum_{\substack{x \leq p \leq 2x \\ p \equiv a \bmod q}} \Psi(u) \\ &= \sum_{\substack{x \leq p \leq 2x \\ p \equiv a \bmod q}} \left(\frac{1}{p} \sum_{\gcd(n,p-1)=1,\, 0 \leq k \leq p-1} \sum \psi\left((\tau^n - u)k\right) \right) \\ &= a_u \sum_{\substack{x \leq p \leq 2x \\ p \equiv a \bmod q}} \frac{1}{p} \sum_{\gcd(n,p-1)=1} 1 \\ &\quad + \sum_{\substack{x \leq p \leq 2x \\ p \equiv a \bmod q}} \frac{1}{p} \sum_{\gcd(n,p-1)=1,\, 0 < k \leq p-1} \sum \psi\left((\tau^n - u)k\right) \\ &= M(x) + E(x), \end{aligned} \tag{32}$$

where $a_u \geq 0$ is a constant depending on the integers $u \neq \pm 1, v^2$ and $q \geq 2$.

The main term $M(x)$ is determined by a finite sum over the trivial additive character $\psi = 1$, and the error term $E(x)$ is determined by a finite sum over the nontrivial additive characters $\psi = e^{i2\pi k/p} \neq 1$.

Take $b > c + 1$. Applying Lemma 5.1 to the main term, and Lemma 4.1 to the error term yield

$$\begin{aligned} 0 &= \sum_{\substack{x \leq p \leq 2x \\ p \equiv a \bmod q}} \Psi(u) \\ &= M(x) + E(x) \\ &= a_u \left(A_q \frac{\operatorname{li}(2x) - \operatorname{li}(x)}{\varphi(q)} \right) + O\left(\frac{x}{\log^b x} \right) + O\left(\frac{1}{\varphi(q)} \frac{x^{1-\varepsilon}}{\log x} \right) \\ &= \delta(u, q, a) \frac{x}{\log x} + O\left(\frac{x}{\log^b} \right), \end{aligned} \tag{33}$$

where $\delta(u, q, a) = a_\ell A_q / \varphi(q) \geq 0$, and $a_u \geq 0$ is a correction factor depending on ℓ.

But $\delta(u, q, a) > 0$ contradicts the hypothesis (31) for all sufficiently large numbers $x \geq x_0$. Ergo, the short interval $[x, 2x]$ contains primes $p = qn + a$ such that the u is a fixed primitive root. Specifically, the counting function is given by

$$\pi_u(x, q, a) = \sum_{\substack{p \leq x \\ p \equiv a \bmod q}} \Psi(u) = \delta(u, q, a) \operatorname{li}(x) + O\left(\frac{x}{\log^b x} \right). \tag{34}$$

This completes the verification. ■.

The determination of the correction factor a_u in a primes counting problem is a complex problem, some cases are discussed in [18], and [22].

The logarithm difference $\operatorname{li}(2x) - \operatorname{li}(x) = x/\log x + O\left(x/\log^2 x\right)$ is evaluated via (28) in Section 5.

7 Special Case $u = 10$

Some of the earliest work on this problem is concerned with the primitive root $u = 10$, see [1], and [4]. A corollary of Theorem 1.1 provides the required density.

Corollary 7.1 *There are infinitely many primes $p \geq 7$ with the primitive root $u = 10$. Moreover, the counting function for these primes satisfies the lower bound*

$$\pi_{10}(x) = \#\{p \leq x : \mathrm{ord}_p(10) = p - 1\} = \delta(10)\frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right), \tag{35}$$

where $\delta(10) = \delta(10, 2, 1) > 0$ is a constant, for all large numbers $x \geq 1$.

Proof: This follows from Theorem 1.1 by replacing the primitive root base $u = 10$. That is,

$$\pi_{10}(x) = \sum_{p \leq x} \Psi(10) = \delta(10)\frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right). \tag{36}$$

Let $u = (sv^2)^k \neq \pm 1$ and $s \equiv 2 \bmod 4$ is squarefree, $k \geq 1$. A formula for computing the density appears in [7, p. 220]. In the decimal case $u = 10$, the density is

$$\begin{aligned} \delta(u) &= \left(1 - \mu(s) \prod_{\substack{p|s \\ p\nmid k}} \frac{1}{p^2 - p - 1} \prod_{\substack{p|s \\ p|k}} \frac{1}{p^2 - p - 1}\right) \prod_{p\nmid k} \left(1 - \frac{1}{p(p-1)}\right) \\ &= \frac{18}{19} \prod_{p \geq 2} \left(1 - \frac{1}{p(p-1)}\right) = 0.357479581392\ldots. \end{aligned} \tag{37}$$

This completes the proof. ■.

References

[1] Bullynck, Maarten, Decimal periods and their tables: a German research topic (1765-1801). Historia Math. 36 (2009), no. 2, 137-160.

[2] Friedlander, John B.; Shparlinski, Igor E. Double exponential sums over thin sets. Proc. Amer. Math. Soc. 129 (2001), no. 6, 1617-1621.

[3] Friedlander, John B.; Hansen, Jan; Shparlinski, Igor E. Character sums with exponential functions. Mathematika 47 (2000), no. 1-2, 75-85 (2002).

[4] Gauss, Carl Friedrich. Disquisitiones arithmeticae. Translated by Arthur A. Clarke. Revised by William C. Waterhouse, Cornelius Greither and A. W. Grootendorst. Springer-Verlag, New York, 1986.

[5] Garaev, M. Z. Double exponential sums related to Diffie-Hellman distributions. Int. Math. Res. Not. 2005, no. 17, 1005-1014.

[6] M. Z. Garaev, A. A. Karatsuba, New estimates of double trigonometric sums with exponential functions, arXiv:math/0504026.

[7] C. Hooley, On Artins conjecture, J. Reine Angew. Math. 225, 209-220, 1967.

[8] Harman, Glyn Prime-detecting sieves. London Mathematical Society Monographs Series, 33. Princeton University Press, Princeton, NJ, 2007.

[9] Hardy, G. H.; Wright, E. M. An introduction to the theory of numbers. Sixth edition. Revised by D. R. Heath-Brown and J. H. Silverman. With a foreword by Andrew Wiles. Oxford University Press, 2008.

[10] Iwaniec, Henryk; Kowalski, Emmanuel. Analytic number theory. AMS Colloquium Publications, 53. American Mathematical Society, Providence, RI, 2004.

[11] Lidl, Rudolf; Niederreiter, Harald. Finite fields. With a foreword by P. M. Cohn. Second edition. Encyclopedia of Mathematics and its Applications, 20. Cambridge University Press, Cambridge, 1997.

[12] Maynard, James. On the Brun-Titchmarsh theorem. Acta Arith. 157 (2013), no. 3, 249-296.

[13] Pieter Moree. Artin's primitive root conjecture -a survey. arXiv:math/0412262.

[14] Moree, Pieter On primes in arithmetic progression having a prescribed primitive root. J. Number Theory 78 (1999), no. 1, 85-98.

[15] Murty, M. Ram; Thangadurai, R. The class number of $Q(\sqrt{-p})$ and digits of $1/p$. Proc. Amer. Math. Soc. 139 (2011), no. 4, 1277-1289.

[16] Montgomery, Hugh L.; Vaughan, Robert C. Multiplicative number theory. I. Classical theory. Cambridge University Press, Cambridge, 2007.

[17] Ribenboim, Paulo, The new book of prime number records, Berlin, New York: Springer-Verlag, 1996.

[18] Stevenhagen, Peter. The correction factor in Artin's primitive root conjecture. Les XXII emes Journees Arithmetiques (Lille, 2001). J. Theor. Nombres Bordeaux 15 (2003), no. 1, 383-391.

[19] Stephens, P. J. An average result for Artin conjecture. Mathematika 16, (1969), 178-188.

[20] Tenenbaum, Gerald. Introduction to analytic and probabilistic number theory. Translated from the Third French edition. American Mathematical Society, Rhode Island, 2015.

[21] Vaughan, R. C. Some applications of Montgomery's sieve. J. Number Theory 5 (1973), 64-79.

[22] Zywina, David. A refinement of Koblitz's conjecture, arXiv:0909.5280.